SUPPLEMENTING NPK WITH ESEFORT ORGANIC GREEN FERTILIZER ON MAIZE, BEANS, RICE AND CASSAVA

| NO. | CROP | NPK AMOUNT | ESEFORT AMOUNT | ACRE | MEASURING ESEFORT NO. OF TIMES TO APPLY | | SPRAYING METHOD | PLANTING METHOD | |
|--|--|-------------------|------------------------------|------|---|---|---|--|--|
| 1 | CASSAVA | 50% (1-2 BAGS) | 250MG DILUTED IN WATER | 1 | 2 CAPS/ KNAPSACK 250MG INTO SIX KNAPSACKS | 1 ST SPRAY: 30 DAYS AFTER PLANTING 2 ND SPRAY: 60 DAYS AFTER 1 ST SPRAY 3 RD SPRAY: 30 DAYS AFTER 3 RD SPRAYING | SPRAY ENOUGH ON LEAVES AND STEMS | (OPTIONAL) SOAK STEMS IN ESEFORT SOLUTION FOR 24 HOURS BEFORE PLANTING | |
| | | | | | | 1 ST SPRAY: | | | |
| | | | | | | 14 DAYS AFTER PLANTING | | SOAK GRAINS IN ESEFORT | |
| 2 | RCICE, MAIZE, BEANS (OTHER GRAINS) | 0% (1-2 BAGS) | 250MG DILUTED IN WATER | 1 | 250MG INTO SIX KNAPSACKS | 2 ND SPRAY: 21 DAYS AFTER 1 ST SPRAYING 3 RD SPRAY: 21 DAYS AFTER 3 RD SPRAYING | SPRAY ENOUGH ON LEAVES AND STEMS | SOLUTION FOR 24 HOURS BEFORE PLANTING | |
| N.B: 1. ESEFORT CONTAINS OVER 70 MICRO-MINERALS 2. ESEFORT FIGHTS AGAINST THE FALL ARMYWORM THAT DESTROYS MAIZE AND OTHER CROPS | | | | | | | | | |
| 3. ESEFORT DESTROYS THE FUNGI THAT DAMAGES LEAVES OF RICE. | | | | | | | | | |
| 4. ESEFORT HELPS TO INCREASES YIELDS. 4. SPRAY IN THE MORNING BEFORE 9:00 AM OR AFTER 4:00PM IN THE EVENING | | | | | | | | | |

SUPPLEMENTING NPK WITH ESEFORT ORGANIC GREEN FERTILIZER ON COCOA, PAWPAW (MOST TREE PLANTS)

| NO. | CROP | NPK AMOUNT | ESEFORT AMOUNT | ACRE | MEASURING ESEFORT | NO. OF TIMES TO APPLY | SPRAYING METHOD | PLANTING METHOD | |
|--|---|---------------------|------------------------------|------|---|---|---|--|--|
| 1 | PAWPAW (PAPAYA) COCOA CHOCOLATE PLANT NURSERY OTHER TREE CROPS | 1.5 BAG | 250MG DILUTED IN WATER | 1 | 2 CAPS/ KNAPSACK 250MG INTO SIX KNAPSACKS | 1 ST SPRAY: AFTER SHOOTING 2 ND SPRAY: 2-3 WEEKS AFTER FIRST SPRAYING | SPRAY ENOUGH ON LEAVES AND STEMS | FOR NURSERY SOAK GRAINS IN ESEFORT SOLUTION FOR 24 HOURS BEFORE PLANTING | |
| 2 | COCOA IMMATURE PLANT COCOA MATURE PLANT | 0% (1-2 BAGS) | 250MG DILUTED IN WATER | 1 | 250MG INTO SIX KNAPSACKS | SPRAY THE IMMATURE PLANT EVERY TWO MONTHS SPRAY PLANT 2-3 WEEKS/ REDUCE 30% OF ORIGINAL AMOUNT OF ESEFORT BUT MAINTAIN THE SAME AMOUNT OF WATER (15) LITRES PER KNAPSACK | SPRAY ENOUGH ON FRUITS LEAVES AND STEMS | | |
| N.B: 1. ESEFORT CONTAINS OVER 70 MICRO-MINERALS 2. ESEFORT EIGHTS AGAINST THE FALL ARMYWORM THAT DESTROYS MAIZE AND OTHER CROPS | | | | | | | | | |

2. ESEFORT FIGHTS AGAINST THE FALL ARMYWORM THAT DESTROYS MAIZE AND OTHER CROPS

SUPPLEMENTING UREA WITH ESEFORT ORGANIC GREEN FERTILIZER ON LEAF VEGETABLE CROPS

| NO. | CROP | UREA | ESEFORT SOLID | ESEFORT LIQUID | ACRE | CONCENTRATION OF ESEFORT | NO. OF TIMES TO APPLY | SPRAYING METHOD |
|---|--|-----------------|------------------|------------------------------|------|---|--|--------------------|
| 1 | CABBAGE, LEAF ONION, SPINACH, CAISIM, LETUCE, CELERY | 50% (1 BAGS) | 1BAG | 250MG DILUTED IN WATER | 1 | 2 CAPS/ KNAPSACK 250MG INTO SIX KNAPSACKS | 15, 25, & 45 (DAY AFTER PLANTING) FOR CABBAGE 15, 25, 35 FOR ONION 1 AND 14 FOR SPINACH 10, 16 FOR CAISIM 15, 25, 35, 35 FOR 15, 25 FOR LETTUCE 20, 30, 40 FOR CELERY | SPRAY ENOUGH |
| N.B: 1. ESEFORT CONTAINS OVER 70 MICRO-MINERALS | | | | | | | | |
| 2. ESEFORT FIGHTS AGAINST THE FALL ARMYWORM THAT DESTROYS MAIZE AND OTHER CROPS | | | | | | | | |
| 3. ESEFORT DESTROYS THE FUNGLI THAT DAMAGES LEAVES OF RICE. 4. ESEFORT HELPS TO INCREASES VIELDS | | | | | | | | |

4. ESEFORT HELPS TO INCREASES YIELDS.

4. SPRAY IN THE MORNING BEFORE 9:00 AM OR AFTER 4:00PM IN THE EVENING